

Relative evaporation probabilities of ^3He and ^4He from the surface of superfluid ^4He

Jonathan P. Warren^a and Charles D.H. Williams^{a,1},

^a*School of Physics, University of Exeter, Stocker Road, Exeter EX4 4QL, United Kingdom*

Abstract

We report a preliminary experiment which demonstrates that ^3He atoms in Andreev states are evaporated by high-energy ($E/k_B \approx 10.2\text{K}$) phonons in a quantum evaporation process similar to that which occurs in pure ^4He . Under conditions of low ^3He coverage, high-energy phonons appear to evaporate ^3He and ^4He atoms with equal probability. However, we have not managed to detect *any* ^3He atoms that have been evaporated by rotons, and conclude that the probability of a roton evaporating a ^3He atom is less than 2% of the probability that it evaporates a ^4He atom.

Keywords: Quantum evaporation; surface; liquid helium; 2-D fermion system

When small quantities of ^3He are added to bulk superfluid ^4He below $T \sim 100\text{mK}$ the atoms occupy so-called Andreev states [1] and form a degenerate two-dimensional fermion system. It has previously been reported that ^3He can be evaporated by phonons [2] in a quantum evaporation [3] process. This report briefly describes an experiment to compare the evaporation probabilities for ^3He and ^4He atoms by positive group-velocity rotons and high-energy phonons.

An electrically heated 1mm^2 thin-film heater in bulk superfluid ^4He was used to generate excitations which travelled ballistically (path length 6.3mm at $\theta = 14^\circ$ to the vertical) to the free surface (Fig. 1). Evaporated atoms were detected with a constant-temperature superconducting bolometer [4] at angle ϕ to the vertical and at radius

5.8mm from the point of intersection of the liquid surface with the centre of the excitation beam.

The high-energy phonons that participate in evaporation have a narrow energy-distribution




Fig. 1. Schematic diagram of the experiment. The bolometer angle ϕ is adjusted with a stepper-motor.

¹ Corresponding author. E-mail: c.d.h.williams@ex.ac.uk

which peaks at $E/k_B = 10.2$ K and cuts off below 10 K [5]. These phonons are generated about a millimeter in front of the heater by a complicated up-scattering mechanism [6]. The ^3He atoms are less tightly-bound to the liquid surface than the ^4He and therefore have a shorter time of flight to the bolometer [2]. The signals were recorded as a function of angle and surface concentration of ^3He . The signal shapes are complicated because the ^3He affects the bolometer responsivity and time-constant, but a simplified analysis is possible by considering only energy-conserving processes involving the dominant 10.2 K phonons, and the relative probabilities of phonon-atom evaporation processes can be inferred as follows.

With an isotopically pure ^4He surface, the measured time τ_4 of the peak in the phonon- ^4He evaporation signal at $\phi = 11^\circ$ (the kinematic angle of evaporation for 10.2 K phonons) is used to establish the arrival time of these phonons at the surface (Fig. 2). Next, the arrival time $\tau_3(\phi)$ of a ^3He atom evaporated by a 10.2 K phonon as a function of evaporation angle ϕ is calculated (inset to Fig. 2). The ^3He atoms have a two-dimensional Fermi distribution of momentum. Those nearest the Fermi energy have the earliest arrival times and travel along paths at the extremes of the angular-distribution of allowed evaporation directions. At a given bolometer position ϕ , the amplitude of the




Fig. 2. A phonon-atom evaporation signal taken at $\phi = 7^\circ$ and $n_{3S} = 1.1 \text{ nm}^{-2}$. The inset shows predicted arrival times associated with evaporation by a 10.2 K phonon.




Fig. 3. Measured fraction of total evaporation signal due to ^3He atoms (points) compared with calculated coverage of surface by ^3He (line).

bolometer signal $S(\phi, t)$ at time $t = \tau_3(\phi)$ is – ignoring minor errors due to imperfect collimation – entirely due to ^3He atoms which have been evaporated by 10.2 K phonons. By integrating the measurements of $S(\phi, \tau_3(\phi))$ and $S(\phi, \tau_4)$ over the solid angle, the relative numbers of ^3He and ^4He atoms evaporated by 10.2 K phonons can be estimated. We conclude that when ^3He coverage is below 0.5 monolayers (3.3 nm^{-2}) a phonon will evaporate an atom of either isotope with equal probability (Fig. 3).

In the light of this result we were surprised to find no evidence, despite a careful search, that positive group-velocity rotons can quantum evaporate ^3He atoms. We believe that the probability that a roton evaporates a ^3He atom is less than our detector noise limit, *i.e.* 2% of its probability of evaporating a ^4He atom.

References

- [1] A.F. Andreev, Soviet Physics JETP **23** 939 (1966).
- [2] M.J. Baird *et al.*, Nature **304** 325 (1983).
- [3] C.D.H. Williams, J. Low Temp. Physics **113** 11 (1998).
- [4] C.D.H. Williams, Meas. Sci. Technol. **1** 322 (1990).
- [5] M.A.H. Tucker and A.F.G. Wyatt, J. Phys. Condens. Matter **6** 2813 & 2825 (1994).
- [6] N. Adamenko *et al.*, Phys. Rev. Lett. **82** 1482 (1999).